

CHARACTERIZATION OF EXTRACELLULAR PROTEASES FROM  
*PSEUDOMONAS* SP. 16D4 AND *DERMACOCCUS* SP. 17D6 ISOLATED FROM  
ARCTIC REGIONS

LIEW KOK JUN

UNIVERSITI TEKNOLOGI MALAYSIA

CHARACTERIZATION OF EXTRACELLULAR PROTEASES FROM  
*PSEUDOMONAS* SP. 16D4 AND *DERMACOCCUS* SP. 17D6 ISOLATED FROM  
ARCTIC REGIONS

LIEW KOK JUN

A dissertation submitted in partial fulfillment  
of the requirements for the award of the  
degree of Master of Science (Biotechnology)

Faculty of Biosciences and Medical Engineering  
Universiti Teknologi Malaysia

JULY 2015

## ACKNOWLEDGEMENT

In preparing this dissertation, I was in contact with many people. They have contributed towards my understanding and thoughts. In particular, I would like to thank my supervisor, Associate Professor Dr. Zaharah binti Ibrahim, for her encouragement, advices and critics. I was greatly appreciate for receiving all the helpful comments and suggestions. I would also like to thank all my senior colleagues and laboratory assistants, especially Dr. Ang and Dr. Ivy, who gave a lot of precious ideas and guidance whenever I encountered difficulties during the research. Without their continued support, this dissertation might not be the same as presented here. In the end, I am grateful to my family, especially my parents, who had giving me their manual supports and advices.

## ABSTRACT

In recent studies, cold-active enzyme has become another subject of interest. It can overcome the limitations of the classical industrial enzymes that normally require higher temperature to function. In this study, 16 bacteria isolated from the Arctic region were screened for their protease activity using Skim Milk agar plate. Two bacteria with lower hydrolysis coefficient (Hc) were selected, they were *Pseudomonas* sp. 16D4 (Hc = 0.339) and *Dermacoccus* sp. 17D6 (Hc = 0.331), and their proteases were produced using “Protease Specific Medium” and were further studied. Both bacteria produced proteases that are associated to growth and the maximal protease activity was reached at 30<sup>th</sup> hour and 24<sup>th</sup> hours of incubation, respectively. Protease produced by *Pseudomonas* sp. 16D4 remained stable at temperature ranging from 0°C to 70°C. Its optimum activity was detected at 0°C, pH 9 or 11 with K<sub>m</sub> value of 1.37 mg/mL and V<sub>max</sub> value of 243.90 Unit/L. As for *Dermacoccus* sp. 17D6, the protease activity remained stable at temperature ranging from 0°C to 40°C and its optimum condition for the assay was at 50°C, pH 7 or 9 with K<sub>m</sub> value of 0.66 mg/mL and V<sub>max</sub> value of 344.83 Unit/L. The SDS-PAGE and zymogram analysis further revealed that the molecular mass of the *Pseudomonas* sp. 16D4’s protease was 40 kDa. Whereas the size for the *Dermacoccus* sp. 17D6’s protease cannot be determined due to the negative result obtained in the zymogram analysis.

## ABSTRAK

Enzim sejuk-aktif telah menjadi satu lagi subjek yang menarik pada zaman ini. Ia dapat mengatasi batasan enzim perindustrian klasik yang biasanya memerlukan suhu yang lebih tinggi untuk berfungsi. Dalam kajian ini, aktiviti protease bagi 16 bakteria dari kawasan Artik telah disaring dengan kegunaan Skim susu agar. Dua bakteria dengan pekali hidrolisis (Hc) yang lebih rendah telah dipilih, mereka adalah *Pseudomonas* sp. 16D4 (Hc = 0.339) dan *Dermaococcus* sp. 17D6 (Hc = 0.331). Dengan kegunaan “Protease Specific Medium”, protease mereka telah dihasilkan dan digunakan dalam kajian seterusnya. Kedua-dua bakteria menghasilkan protease yang berkaitan dengan pertumbuhan dan maksimum protease aktiviti boleh dicapai pada jam 30 dan jam 24, masing-masing. Protease yang dihasilkan oleh *Pseudomonas* sp. 16D4 dapat mengekalkan kestabilannya dari suhu 0°C hingga 70°C. Aktiviti optimumnya pula dikesan pada 0°C, pH 9 atau 11 dengan nilai  $K_m = 1.37$  mg/mL dan nilai  $V_{max} = 243,90$  Unit / L. Bagi *Dermaococcus* sp. 17D6 pula, enzimnya dapat mengekalkan kestabilan pada suhu antara 0°C hingga 40°C dan keadaan optimum untuk enzim itu adalah 50°C, pH 7 atau 9 dengan nilai  $K_m = 0.66$  mg mL dan nilai  $V_{max} = 344,83$  Unit / L. Analisis SDS-PAGE dan zymogram telah mendedahkan bahawa molekul saiz bagi enzim *Pseudomonas* sp. 16D4 adalah 40 kDa. Bagi enzim *Dermaococcus* sp. 17D6 pula, saiznya tidak boleh dianggarkan kerana zymogram analisis tidak memberi sebarang data.